Environmental Biology of Fishes **46**: 417–419, 1996.

Volume 46

Issue 1, pp. 1–108: May 1996
Issue 2, pp. 109–218: June 1996
Issue 3, pp. 219–328: July 1996
Issue 4, pp. 329–422: August 1996

Volume contents

Invited editorial

Why do fish have so few roundworm (nematode) parasites, by R.C. Anderson 1–5
Threatened fishes of the world: Meda fulgida *Girard*, 1856 (*Cyprinidae*), *by P.C. Marsh* 6

Full papers

Does body and fin form affect the maneuverability of fish traversing vertical and horizontal slits? by P.W. Webb, G.D. LaLiberte & A.J. Schrank 7–14
Habitat correlates of the distribution and biomass of Seychelles' reef fishes, by S. Jennings, D.P. Boullé & N.V.C. Polunin 15–25
Threatened fishes of the world: Clinus spatulatus *Bennett, 1983 (Clinidae), by A.K. Whitfield* ... 26
Intertidal territoriality and time-budget of the surgeonfish, *Acanthurus lineatus*, in American Samoa, by P. Craig 27–36
Microhabitat partitioning in a diverse assemblage of darters in the Allegheny River system, by J.R. Stauffer, Jr., J.M. Boltz, K.A. Kellogg & E.S. van Snik 37–44
Development and utilization of a population growth history of Red Lake walleye, *Stizostedion vitreum*, by M.J. Cyterski & G.R. Spangler 45–49
Threatened fishes of the world: Prototroctes oxyrhynchus *Günther, 1870* (*Prototroctidae*), by R.M. McDowall 60
In situ determination of food daily ration in fish: review and field evaluation, by D. Héroux & P. Magnan 61–74
Effects of cadmium on foraging behaviour of lake charr, *Salvelinus namaycush*, by M. Kislalioglu, E. Scherer & R.E. McNicol 75–82
Feeding of ecology of three *Astyanax* species (Characidae, Tetragonopterinae) from a floodplain lake of Mogi-Guaçú River, Paraná River Basin, Brazil, by K. Eichbaum Esteves 83–101

Brief communication

The influence of parental care behaviour on localized nest spacing in smallmouth bass, *Micropterus dolomieu*, by R.J. Scott, ... 103–107

Full papers

Visual constraints on migration behavior of juvenile French grunts, by W.N. McFarland & C.M. Wahl 109–122
Variation in reef associated assemblages of the Lutjanidae and Lethrinidae at different distances offshore in the central Great Barrier Reef, by S.J. Newman & D.McB. Williams 123–138

Aspects of the reproductive biology of the concha wrasse, *Nelabrichthys ornatus*, at Tristan da Cunha, by T.G. Andrew, C.D. Buxton & T. Hecht 139–149
Threatened fishes of the world: Nannoperca oxleyana *Whitley, 1940 (Nannopercidae), by A.H. Arthington & C.J. Marshall* 150
Ecological fish production in the inland delta of the Middle Danube, a floodplain river, by J. Holčík 151–165
Threatened fishes of the world: Salvethymus svetovidovi *Chereshnev & Scopets, 1990 (Salmonidae), by I.A. Chereshnev* 166
Trophically-based assessment of the impacts of deepwater sewage disposal on a demersal fish community, by N.M. Otway, D.J. Sullings & N.W. Lenehan 167–183
Threatened fishes of the world: Hypseleotris dayi *Smith, 1950 (Eleotridae), by M.N. Bruton* 184
Why do juvenile fourhorn sculpin, *Triglopsis quadricornis*, appear in the pelagic habitat at night? by J. Hammar, E. Bergstrand & O. Enderlein 185–195
Growth, survival, and starvation resistance of Colorado squawfish larvae, by K.R. Bestgen 197–209

Brief communication
Relative effects of turbidity and light intensity on reactive distance and feeding of an estuarine fish, by M.C. Benfield & T.J. Minello 211–216

Book critique
Half a journey, by D.W. Larson 217–218

Invited editorial
The whale shark, *Rhincodon typus*, is a livebearer: 300 embryos found in one 'megamamma' supreme, by S.-J. Joung, C.-T. Chen, E. Clark, S. Uchida & W.Y.P. Huang 219–223

Full papers
The life history of the tropical shad *Tenualosa toli* from Sarawak: first evidence of protandry in the Clupeiformes? by S.J.M. Blaber, D.A. Milton, J. Pang, P. Wong, O. Boon-Teck, L. Nyigo & D. Lubim 225–242
Persistence of egg pocket architecture in redds of chum salmon, *Oncorhynchus keta*, by N.P. Peterson & T.P. Quinn 243–253
Threatened fishes of the world: Eucyclogobius newberryi *Girard, 1857 (Gobiidae), by K.D. Lafferty, R.O. Swenson & C.C. Swift* 254
Ontogenetic habitat shifts by *Galaxias gracilis* (Galaxiidae) between the littoral and limnetic zones of Lake Kanono, New Zealand, by D.K. Rowe & B.L. Chisnall 255–264
Variation in abundance and survival of fish larvae in shallow eutrophic lake Tjeukemeer, by W.M. Mooij 265–279
Threatened fishes of the world: Galaxias gracilis *McDowall, 1967 (Galaxiidae), by R.M. McDowall & D.K. Rowe* 280
Both-ways sex change in monogamous coral gobies, *Gobiodon* spp., by Y. Nakashima, T. Kuwamura & Y. Yogo 281–288
The effect of group size on space use and aggression at a concentrated food source in blue gouramis, *Trichogaster trichopterus* (Pisces: Belontiidae), by S. Syarifuddin & D.L. Kramer 289–296
Feeding styles of five species of bottom-feeding fishes of the high Paraná River, by R. Fugi, N.S. Hahn & A.A. Agostinho 297–307

Threatened fishes of the world: Pseudobarbus quathlambae *Barnard, 1938) (Cyprinidae), by J.A. Cambray* 308
Critique of proportional hypotheses and methods for back-calculation of fish growth, by M. Živkov 309–320

Brief communication
On the mechanism of sperm release in three gobiid fishes (Teleostei: Gobiidae), by A. Marconato, M.B. Rasotto & C. Mazzoldi 321–327

Full papers
A quantitative description of the reproductive biology of the Mediterranean blenny *Aidablennius sphynx* (Teleostei, Blenniidae) in its natural habitat, by S.B.M. Kraak 329–342
Effect of density on the reproductive behaviour of the territorial male demoiselle *Chromis dispilus* (Pisces: Pomacentridae), by C.W. Barnett & N.W. Pankhurst 343–349
Threatened fishes of the world: Maccullochella ikei *Rowland, 1985 (Percichthyidae), by S.J. Rowland* 350
Comparison of fish assemblages among an artificial reef, a natural reef and a sandy-mud bottom site on the shelf off Iwate, northern Japan, by T. Fujita, D. Kitagawa, Y. Okuyama, Y. Jin, Y. Ishito & T. Inada 351–364
Evidence for ram suspension feeding by the piscivore, *Seriola dumerili* (Carangidae), by S.L. Sanderson, C.R. Chesnutt & P.S. Lobel 365–373
Threatened fishes of the world: Iranocypris typhlops *Bruun & Kaiser, 1994 (Cyprinidae), by B.W. Coad* 374
Selective ingestion of detritus by a north temperate omnivorous fish, the juvenile white sucker, *Catostomus commersoni*, by M.O. Ahlgren 375–381
Threatened fishes of the world: Notropis simus pecosensis *Gilbert & Chernoff, 1982 (Cyprinidae), by F.B. Furlow* 382
Spatial and annual variation in the diets of juvenile smallmouth bass, *Micropterus dolomieu*, by R.S. Easton, D.J. Orth & J.R. Voshell, Jr. 383–392
Effect of stream microhabitat characteristics on rate of net energy gain by juvenile smallmouth bass, *Micropterus dolomieu*, by M.J. Sabo, D.J. Orth & E.J. Pert 393–403

Brief communications
Parallel swims as a means of intra- and interspecific assessment in stream fishes, by C.E. Johnston, J.W. Armbruster & C.A. Laird 405–408
Clutch and egg size variation in the banded darter, *Etheostoma zonale*, from three sites in Arkansas, by J.M. Guill & D.C. Heins 409–413

Authors' index 415

Acknowledgements for refereeing 416

Volume contents 417–419

Instructions to authors 421–422